

Packing problems are often very interesting. In this problem we will try to put five L's in a circle. But before that we will try to define L-shape for this problem.

We can divide a  $2a \times 2a$  square into four  $a \times a$  squares as shown in figure 1. Now if we remove one of those smaller squares from the figure, the new shape we get is referred as L-shape or L in this problem. The length of the smaller sides of this shape is  $a$ .




Figure 1: Defining L




Figure 2: Packing 5 identical L's in a circle of minimum radius

Figure 2 shows how five identical Ls can be put into a circle of minimum possible radius (This is found by David Cantrell but not formally proved). Given the length of the smaller side  $a$  of an L, your job is to find the minimum possible radius of the circle in which five such L's can be put in the fashion shown in figure 2.

### Input

Input file contains at most 5000 lines of inputs. Each line contains a floating-point number  $a$  ( $0 < a < 10000$ ). The meaning of  $a$  is given in the problem statement. This floating-point numbers should have 7 digits after the decimal point. Input is terminated by a line containing  $a - 1$ . This line should not be processed.

### Output

For each line of input produce one line of output. This line contains a floating-point number  $R$  that denotes the Minimum possible radius of the desired circle. This value should have 10 digits after the decimal point. Errors less than  $\max(10^{-9}, 10^{-9} * RJ)$  will be ignored. Here  $RJ$  is the value of  $R$  produced by judge solutions.

### Sample Input

```
0.0000010
0.0000001
-1
```

### Sample Output

```
0.0000026406
0.000002641
```