

You are given an array of  $N$  integers:  $A_1, A_2, \dots, A_N$ . You have to process  $Q$  queries on this array, where a query will be a pair of integers  $(L, R)$ .

For each query, you have to find the count of Divisor-free numbers in the number sequence  $S$ , where  $S = A_L, A_{L+1}, \dots, A_R$ . A number  $A_i$  from the sequence  $S$  will be called Divisor-free if there is no  $A_j$  ( $i \neq j$ ) in  $S$  such that  $A_j$  is a divisor  $A_i$ .

## Input

The first line of the input contains an integer  $T$  ( $T \leq 5$ ) denoting the number of test cases. The first line of each test case contains two integers  $N$  and  $Q$  ( $1 \leq N, Q \leq 10^5$ ). The following line contains  $N$  space separated integers  $A_1, A_2, \dots, A_N$  where  $1 \leq A_i \leq 10^6$ . In each of the next  $Q$  lines, there will be two integers  $(L, R)$  representing a query ( $1 \leq L \leq R \leq N$ ).

## Output

For each test case, print the case number in the format 'Case  $X$ :' (here,  $X$  is the serial of the test case). Then print  $Q$  lines containing the answer for each query.

## Sample Input

```
2
10 5
4 6 2 7 5 11 14 21 13 2
2 6
4 8
2 8
3 7
4 9
5 3
4 6 8 1 5
1 5
2 3
3 3
```

## Sample Output

```
Case 1:
4
3
4
4
4
Case 2:
1
2
1
```